

Mathematical Concepts (G6012)



Lecture 13

Martin Ritchie

FUNCTIONS

(Maths) Functions


A function is like a vending machine:



Elements of the **domain** of the function

The elements of the **co-domain (image)** of the function

Notation



Functions are also often called **map**, **mapping** or **transformation**.

Examples: Polynomials

$$f : \mathbb{R} \rightarrow \mathbb{R}_+$$

$$x \mapsto x^2$$

Alternative
notation:

$$f(x) = x^2$$

This means ...

$$1 \mapsto 1$$

$$f(1) = 1$$

$$\sqrt{2} \mapsto 2$$

$$f(\sqrt{2}) = 2$$

$$2 \mapsto 4$$

$$f(2) = 4$$

$$-3 \mapsto 9$$

$$f(-3) = 9$$

$$4 \mapsto 16$$

$$f(4) = 16$$

$$5 \mapsto \dots$$

$$f(5) = \dots$$

More general

A **polynomial** is a function of the form

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$f(x) = \sum_{n=0}^k a_n x^n$$

degree of the polynomial

coefficients

Example: Matrices (more later)

$A \in \mathcal{M}(m, n)$ i.e. A is a $m \times n$ matrix

We have seen before that A describes a function (transformation) by:

$$A : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$\vec{x} \mapsto A \cdot \vec{x}$$

Matrix-vector multiplication

Examples: “Not” function

$$\begin{aligned} \textit{not} : \{0, 1\} &\rightarrow \{0, 1\} \\ x &\mapsto \textit{not}(x) \end{aligned}$$

x	not(x)
0	1
1	0

Alternative
notation:

$$0 \mapsto 1$$

$$1 \mapsto 0$$

Alternative 2:

$$\textit{not}(0) = 1$$

$$\textit{not}(1) = 0$$

Examples: “+” function

$$+ : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$$

$$(x, y) \mapsto x + y$$

This means:

$$(2, 2) \mapsto 2 + 2 = 4$$

$$(2, -3) \mapsto 2 + (-3) = 2 - 3 = -1$$

Alternative notation:

$$+(1, 4) = 5$$


Please note ...

Functions are **single valued**, i.e. for any number x from the domain of the function there is **one unique number** $y = f(x)$ from the co-domain.

BB Graph of a function


The **graph** of a function are all pairs (points) $(x, f(x))$

$$f(x) = 1.5 \cdot x$$



A linear function

Warning: Why is this **not** a graph of a function?








What about these ones?

Further Clinical results from Dr. Alan Fermaine's Trials


Incredible Speed of Action

Penis Dilatation %




Some example graphs of functions

$$f(x) = x^2$$



A parabola

$$f(x) = \exp(x) = e^x$$




The exponential function

Other functions you should have seen/know

$$f(x) = \sin(x)$$

$$f(x) = \cos(x)$$



LIMITS

The idea

- Problem: **Sequence** of numbers, e.g.

$$a_n = \frac{1}{n} \text{ and } n = 1, 2, \dots$$

What happens to $\frac{1}{n}$ if we make n ever larger?

Mathematicians write $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$

Two *typical* cases:

The numbers go ever larger (the series **diverges**),
or (more interesting) it **converges** to a number
(here 0).

Formal Definition


A sequence a_n converges to a number $x \in \mathbb{R}$
if for all (small) $\epsilon > 0$ there is a number $N \in \mathbb{N}$
such that $|a_n - x| < \epsilon$ for all $n \geq N$.

Intuition: **BB**

BB Intuition: Convergence

Blue dots: a_n up to some $N \in \mathbb{N}$

Green dots: a_n for all larger n



Examples

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0 \quad \mathbf{BB}$$

The limit for the sequence $a_n = n$ for n to infinity does not exist, it grows without bound, people write:

$$\lim_{n \rightarrow \infty} n = \infty$$

BB

Claim: $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$

Proof: Let $\epsilon > 0$.

We choose $N = \text{floor} \left(\frac{1}{\epsilon} \right) + 1$.

Then, for all $n \geq N$, we know

$$\left| \frac{1}{n} - 0 \right| \leq \left| \frac{1}{N} \right| < \left| \frac{1}{\frac{1}{\epsilon}} \right| = \epsilon$$

Practical rules for limits

- If all limits involved exist,

$$\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$$

$$\lim_{n \rightarrow \infty} (a_n \cdot b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$$

$$\lim_{n \rightarrow \infty} (k \cdot a_n) = k \cdot \lim_{n \rightarrow \infty} a_n$$

“Thumb” rules for calculating limits

(Please don't show this slide to proper
mathematicians)

$$\text{If } \lim_{x \rightarrow \infty} x = \infty$$

$$\text{then } \lim_{x \rightarrow \infty} \frac{1}{x} = \frac{1}{\infty} = 0$$

Example

$$f(x) = \exp(-x) = e^{-x}$$

$$\lim_{x \rightarrow \infty} \exp(-x) = \lim_{x \rightarrow \infty} \frac{1}{\underbrace{\exp(x)}_{\rightarrow \infty}} = 0$$

Some more

(Please don't show this slide to proper mathematicians)

$$k \cdot \infty = \begin{cases} \infty & \text{if } k > 0 \\ -\infty & \text{if } k < 0 \end{cases}$$

$$\frac{k}{0} = \begin{cases} \infty & \text{if } k > 0 \\ -\infty & \text{if } k < 0 \end{cases}$$

Another example

(Please don't show this slide to proper mathematicians)

$$\lim_{x \rightarrow 0} \exp\left(-\frac{1}{x}\right) =$$
$$\exp\left(-\frac{1}{0}\right) = \exp(-\infty) = \frac{1}{\exp(\infty)} = 0$$

Warning: Not everything converges or diverges (goes to ∞)

What about $\lim_{n \rightarrow \infty} \sin(n)$? **It does not exist at all!**

Big O notation: Speed of divergence

- Is used especially in Computer Science
- A function f is $\mathcal{O}(g)$, where g is another function, if $|f(x)| \leq k \cdot |g(x)|$ for $x \rightarrow \infty$ and some $k \in \mathbb{R}$.
- In other words, if f is **on the order of** g for large x (or less than g ...)

Examples

$$f(x) = x^2 + 5x + 10 \text{ is } \mathcal{O}(x^2)$$

It is also $\mathcal{O}(x^3)$

$$x\sqrt{x} \text{ is } \mathcal{O}(x^2)$$


Note: if f is $\mathcal{O}(g)$ and g converges to 0, then f converges to 0 as well!

DERIVATIVES

Slope, Derivative

First for **linear functions**:

$$f(x) = a \cdot x$$



The slope or derivative is the ratio of the change of $f(x)$ and the change of x .


$$\frac{\Delta y}{\Delta x} = \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

BB

BB

$$\frac{\Delta y}{\Delta x} = \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

$$f(x) = 1.5 \cdot x$$



BB Calculating the linear example

$$f(x) = 1.5 \cdot x$$

$$\frac{\Delta y}{\Delta x} = \frac{f(x + \Delta x) - f(x)}{\Delta x}$$


$$= \frac{1.5 \cdot (x + \Delta x) - 1.5 \cdot x}{\Delta x} = 1.5$$


Linear functions are easy ...


- Because the slope is the same everywhere Δx
- We can make any size we want and get the same value


Generalisation for any smooth function


- Locally **any** smooth function looks more and more linear the further we zoom in:











Derivative of a smooth function

- The derivative of a smooth function is the value the ratio $\frac{f(x + \Delta x) - f(x)}{\Delta x}$ converges to for smaller and smaller Δx , mathematicians write

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

In the form as for the limits
before

Sequence $a_n = \frac{f(x + \frac{1}{n}) - f(x)}{\frac{1}{n}}$

$$f'(x) = \lim_{n \rightarrow \infty} \frac{f(x + \frac{1}{n}) - f(x)}{\frac{1}{n}}$$

Alternative notations

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function
 $x \mapsto f(x)$

Then the derivative of f is denoted as

$$f'(x) = \frac{df(x)}{dx} = \frac{df}{dx} = \frac{d}{dx}f$$

Note ...

The derivative $f'(x)$ of a function is again a function because we can calculate it for any point x .

BB

Example - Derivative of $f(x) = x^2$

$$\begin{aligned}\frac{d}{dx}x^2 &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^2 + 2x \Delta x + (\Delta x)^2 - x^2}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} 2x + \Delta x = 2x\end{aligned}$$

Applications

- If $f(x)$ is your **distance** from home as a function of the time x . Then $f'(x)$ is the **speed** you are driving towards (or away from) home.
- If you take the derivative of the derivative $f''(x)$, that would be your **acceleration**.

(These are important for animating things!)

More Applications

- If $f(x)$ describes the **height** of a hill, then $f'(x)$ is the **steepness**.
- $f(x)$ is your **total money** as a function of time, $f'(x)$ is your instantaneous **spending rate**.
- (your example here)

Derivative of a polynomial

- Last time:

$$f(x) = ax \quad \text{then} \quad f'(x) = \frac{d}{dx} f(x) = a$$

- For $f(x) = x^2$ we saw just now $f'(x) = 2x$

- Generally, for $f(x) = x^n$
the derivative is $f'(x) = nx^{n-1}$

For those interested: General case

$$f(x) = x^n$$

$$\begin{aligned}\frac{d}{dx}x^n &= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^n - x^n}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{x^n + nx^{n-1}\Delta x + \mathcal{O}((\Delta x)^2) - x^n}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} nx^{n-1} + \mathcal{O}(\Delta x) = nx^{n-1}\end{aligned}$$

Derivatives: Basic rules

Rule name	Function	Derivative
Polynomials	$f(x) = x^n$	$f'(x) = n x^{n-1}$
Constant factor	$g(x) = a f(x)$	$g'(x) = a f'(x)$
Sum and Difference	$h(x) = f(x) + g(x)$	$h'(x) = f'(x) + g'(x)$

Examples: Polynomial rule

Example 1

$$f(x) = \sqrt{x} = x^{\frac{1}{2}}$$

$$f'(x) = \frac{1}{2}x^{-\frac{1}{2}} = \frac{1}{2} \frac{1}{\sqrt{x}}$$

Example 2

$$g(x) = \frac{1}{x^n} = x^{-n}$$

$$g'(x) = -n x^{-n-1} = \frac{-n}{x^{n+1}}$$

Special functions

Function	Derivative
$\exp(x) = e^x$	$\exp(x) = e^x$
$\log(x) = \ln(x)$	$\frac{1}{x}$
$\sin(x)$	$\cos(x)$
$\cos(x)$	$-\sin(x)$